

LA-UR-20-23926

Approved for public release; distribution is unlimited.

Title: Biogenic uranium isotope fractionation

Author(s): Marti-Arbona, Ricardo
Jemison, Noah
Williams, Robert F.
Boukhalfa, Hakim
Yeager, Chris Michael
Xu, Ning
Vesselinov, Velimir Valentinov

Intended for: Share with collaborator

Issued: 2020-05-28

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Biogenic uranium isotope fractionation

Team presentation

**Robert Williams (PI, B-11) Hakim
Boukhalfa (Co-PI, EES-14)
Ricardo Martí-Arbona (Co-PI, B-11)
Co-I's
Chris Yeager (C-CDE)
Ning Xu (C-AAC)
Velimir V. Vesselinov (EES-16)
Noah Jemison (former EES-14)**


Outline

- ▶ Background and relevance
- ▶ Project objectives
- ▶ Tasks and timeline
- ▶ Current Activities
 - ▶ Task 1: Controlled culture growth
 - ▶ Task 2: Enzymatic reduction of uranium oxides
 - ▶ Task 3: Cellular location of uranium reduction and precipitation
- ▶ Future work
- ▶ Achievements

Uranium biogeochemistry


- ▶ Uranium chemistry controls how U ore deposits form, how to clean up U contamination, and how U is distributed in the environment
- ▶ U is redox-active
 - ▶ U(VI)- soluble and mobile in water systems
 - ▶ U(IV)- relatively insoluble and primarily in sediment systems
- ▶ Common method to remove U from solution is microbial U(VI) reduction
- ▶ However, uranium concentrations impacted by multiple chemical and physical processes

a) June 1994



Uranium isotopes

- ▶ Uranium isotopes ($^{238}\text{U}/^{235}\text{U}$) can provide a more direct indicator of U(VI) reduction
- ▶ Not strongly affected by adsorption or physical processes
- ▶ Microbial U(VI) reduction- preferential reduction of ^{238}U (Basu et al., 2014; Stylo et al., 2015)
 - ▶ Less ^{238}U remaining in U(VI) solution
- ▶ Abiotic reduction produces a large range of isotopic fractionation (Brown et al., 2018; Stylo et al., 2015)
- ▶ What are the processes and mechanisms controlling isotope fractionation?




Basu et al., 2014

What affects fractionation?

- ▶ Isotopic fractionation defined as:

$$\epsilon = \frac{{}^{238}\text{U}/{}^{235}\text{U}_{\text{U(IV) product}}}{{}^{238}\text{U}/{}^{235}\text{U}_{\text{U(VI) reactant}}}$$
- ▶ Aqueous chemistry impacts isotope fractionation during abiotic experiments
- ▶ May be due to U(VI) speciation or reduction rate
- ▶ U(VI) reduction to U(V) and then disproportionation to U(IV) and U(VI)?
- ▶ Biosorption, bioaccumulation, and bioreduction mechanisms?
- ▶ What are the primary factors controlling the magnitude and direction of isotope fractionation during U(VI) reduction?
- ▶ Can we reliably apply ${}^{238}\text{U}/{}^{235}\text{U}$ to track and quantify U(VI) reduction in natural environments?



Project objectives

- ▶ The underlying goal is to determine the mechanistic driver(s) of U fractionation, probing processes from the initial interaction between the cell and soluble U to the accumulation of U mineral precipitates near or within the cell.
- ▶ We will focus on the characterization of three aspects of uranyl bioreduction that likely control U isotope fractionation:
 - ▶ **1.** kinetic controls that dictate U adsorption, sequestration, and/or uptake and its subsequent reduction;
 - ▶ **2.** cellular processes that support the electron transport pathways and enzymatic reduction of uranium;
 - ▶ **3.** characterization and mapping of the cellular location of U reduction and precipitation.

Impact of research

- ▶ Determining the primary mechanisms of U isotope fractionation would establish LANL as a leader in environmental isotope measurements
- ▶ Develop our capabilities for tracing environmental biogeochemical reactions
- ▶ Gain more recognition for emerging isotope measurements and applications

100

Schedule and Milestones

[illegible]

FY-19 activities

- ▶ Task 1: Controlled culture growth
 - ▶ Strain selection and initial cultivation
 - ▶ Parameter sensitivity screening and statistical analysis
 - ▶ Cultivation experiments with reduced number of parameters
 - ▶ Nano SIM, TEM, SEM characterization
 - ▶ XAS characterization of select samples
- ▶ Task 2: Enzymatic reduction of uranium oxides
 - ▶ Protein Expression and Purification
 - ▶ Activity Characterization
- ▶ Task 3: Cellular location of uranium
 - ▶ Uranium uptake
 - ▶ Cellular Sorption
 - ▶ Intracellular uptake

Strain selection and initial cultivation

- ▶ Currently, growing *Shewanella oneidensis* and *Pelosinus* strain UFO1
 - ▶ Both capable of U(VI) reduction
 - ▶ *Shewanella*- facultative anaerobe, gram (-)
 - ▶ *Pelosinus*- strict anaerobe, gram (+)
- ▶ Any differences in isotope fractionation due to different microbial mechanisms?
- ▶ How does aqueous chemistry affect isotope fractionation during reduction by these microbes?
- ▶ Could microbial uptake impact observed isotope fractionation?

Parameter sensitivity screening

▶ **Abiotic U(VI) reduction experiments**

- ▶ Abiotic experiments eliminate some of the complexity of microbial experiments
- ▶ These experiments allow us to screen for what parameters strongly affect isotope fractionation during U(VI) reduction
- ▶ How does U(VI) speciation, solution chemistry, and reduction rate impact isotope fractionation?


Methods

- ▶ Performed batch experiments in an anaerobic chamber where reductant was added to U solutions
- ▶ Reductants: FeS and Na₂S with quartz
- ▶ Varied chemistry
 - ▶ pH (6.5 and 7)
 - ▶ bicarbonate (6 and 2mM)
 - ▶ Ca (0, 1, and 2mM)
 - ▶ Mg (0 and 10mM)
 - ▶ MOPS pH buffer (10 and 70mM)
- ▶ Calculated U(VI) speciation and adsorption coefficients (K_D) using CrunchTope

Experiment	mM Ca	mM Mg	mM HCO ₃	mM MOPS	pH	reductant
1	1	0	6	70	7	FeS
2	1	0	6	70	7	FeS
3	0	0	6	70	7	FeS
4	0	0	6	70	7	FeS
5	2	0	6	70	7	FeS
6	2	0	6	70	7	FeS
7	2	0	6	70	7	FeS
8	0	10	6	70	7	FeS
9	1	0	6	10	7	FeS
10	1	0	2	70	7	FeS
11	1	0	6	70	6.5	FeS
12	1	0	6	70	6.5	FeS
13	1	0	6	70	7	HS
14	1	0	2	70	7	HS


Isotope methods

- ▶ Collected samples over time as U(VI) was reduced
- ▶ Samples filtered to remove reductant and U(IV)
- ▶ Analyzed remaining U(VI)
- ▶ Added ^{233}U - ^{236}U double spike to U(VI) samples to account for mass bias
- ▶ Measured U(VI) concentrations and $\delta^{238}\text{U}$ on a multi-collector ICPMS (MC-ICPMS) at University of Illinois
- ▶ $\delta^{238}\text{U} = \left(^{238}\text{U}/^{235}\text{U}_{\text{sample}} / ^{238}\text{U}/^{235}\text{U}_{112\text{A std}} - 1 \right) * 1000\text{‰}$




Concentration data

2mM Ca




pH 6.5




Isotope Data


2mM Ca




pH 6.5



Fractionation mechanisms?



Fractionation mechanisms?



Parameter sensitivity discussion

- ▶ Adsorption coefficient much more strongly correlated with ε than U(VI) speciation or reduction rate
- ▶ Aqueous chemistry can influence U isotope fractionation through adsorption and then reduction of U(VI)
- ▶ U(VI) adsorption induces an isotopic fractionation of $\sim 0.2\text{‰}$ (adsorbed U(VI) enriched in ^{235}U) (Jemison et al., 2016)
- ▶ Fractionation remains intact when K_D is high, but with low K_D , ε is dominated by U(VI) reduction (^{238}U preferentially reduced)

Parameter sensitivity discussion

- ▶ With high K_D , most U(VI) that is adsorbed is reduced
- ▶ With low K_D , more U(VI) can desorb and communicate with the aqueous U(VI) pool
- ▶ What about microbial U(VI) reduction?
 - ▶ Need to test how aqueous chemistry impacts U isotope fractionation

Enzymatic reduction of uranium oxides

- ▶ Chris will update
 - ▶ **Protein Expression and Purification**
 - ▶ **Activity Characterization**

FY-19 Accomplishments

- ▶ Jemison, N.; Reimus, P.; Harris, R.; Boukhalfa, H.; Clay, J.; Chamberlain, K. Reduction and potential remediation of U(VI) by dithionite at an in-situ recovery mine: insights gained by $\delta^{238}\text{U}$. Applied Geochemistry. Submitted. (LA-UR-19-27182)
- ▶ Jemison, N.; Boukhalfa, H.; Marti-Arbona, R.; Yeager, C.; Ning, X. Mechanisms of Uranium Isotope Fractionation. Poster, Goldschmidt 2019. (LA-UR-19-22703)

Current activities

- ▶ Task 1: Controlled culture growth
 - ▶ Strain selection and initial cultivation
 - ▶ Parameter sensitivity screening and statistical analysis
 - ▶ Cultivation experiments with reduced number of parameters
 - ▶ Nano SIM, TEM, SEM characterization
 - ▶ XAS characterization of select samples
- ▶ Task 2: Enzymatic reduction of uranium oxides
 - ▶ Protein Expression and Purification
 - ▶ Activity Characterization
- ▶ Task 3: Cellular location of uranium
 - ▶ Uranium uptake
 - ▶ Cellular Sorption
 - ▶ Intracellular uptake

Microbial Pu reduction

- ▶ Will soon reduce Pu(VI) to Pu(V) and Pu(IV) using *Shewanella* and *Pelosinus*
- ▶ Allows us to see fractionation for each electron step
 - ▶ U(VI) reduction does not produce significant amounts of stable U(V) species
- ▶ First study on Pu isotope fractionation during natural reduction processes



External Collaborators

- ▶ Tom Johnson (University of Illinois- Urbana-Champaign)
- ▶ John Cliff (Pacific Northwest National Laboratory)
- ▶ ??